


### METHOD (complete one survey form per pond)

**Aims:** To find and map Bearded Stonewort on the sites where it occurs and record a measure of abundance at each pond. The aim is to find out whether Bearded Stonewort is i) present, ii) get an approximate idea of its location and abundance, iii) collect physical data that can be used to assess the reasons for any change recorded on future visits, and iv) look in any adjacent ponds to see if Bearded Stonewort is present or absent.

- Equipment: It's helpful to take a camera (e.g. mobile phone camera) to take confirmatory photos of Bearded Stonewort, to take photos of your survey pond for the record, and to take a photograph of your sketch maps if you don't have access to a scanner alternatively you can post your survey forms to Freshwater Habitats Trust.
- Survey timing: Bearded Stonewort is best surveyed in the autumn, between September and October.
- Where to look: Bearded Stonewort typically grows in shallow pools and may be recorded anywhere in the pond, including in the drawdown zone the area that is wet in winter, but progressively dries out in summer. Search for it across all of the pond's dry marginal areas and in shallow water.
- Survey the area indicated on your map: The pond may have a previous record for Bearded Stonewort, although the plant may not have been recorded for some time, or it may be a new site. Search the area indicated in your site pack for Bearded Stonewort plants, and if found, <u>estimate abundance</u> (see below), draw a sketch map to show <u>the location and extent of Bearded Stonewort</u> and, <u>fill out the habitat survey form overleaf</u>.
- How to estimate abundance: Bearded Stonewort grows as a dense cluster, so it can be difficult to count individual plants. Abundance therefore needs to be an estimate of plant cover. To help standardised these estimates we are using two measures of abundance, the area in m<sup>2</sup> and the percentage of the pond occupied.

**Measurement 1.** <u>Area covered by Bearded Stonewort</u>: The aim is to record the total *area* of the Bearded Stonewort (in  $m^2$ ). To do this, record the size of each patch of plants, e.g.  $(1m \times 1m) + (1m \times 2m) = 3m^2$ . It can help to record a number of patches by imagining them grouped together to make a square or rectangle. **Note: We only need to know the total area of Bearded Stonewort to monitor the pond**, but the space overleaf can help you to add up the different patches.








Patch = 2m<sup>2</sup>

Bearded Stonewort may occur at different **densities** in each patch: sometimes growing close together, and at other sites more widely separated. You need to *standardise the density*. To do this imagine more sparsely growing plants are pushed together to grow at their maximum *natural* density (see photo below).

### Measurement 2. Percentage of the pond occupied by Bearded Stonewort:

The aim is to estimate the percentage of the pond that Bearded Stonewort occupies. Use the density chart (right), or imagine that the plants are grouped together at their maximum *natural* density in one part of the pond.



1m

- Recording absence: If Bearded Stonewort <u>is not</u> found, please record this, and continue to fill out the environmental sheet. The findings will help identify reasons for the plant's absence.
- Check other likely habitats in the surrounds: Finding out if Bearded Stonewort occurs in other likely habitats within the same site helps us to understand if the species is part of a larger population, which may be important for its survival. Visit other likely habitat patches within the site to see if Bearded Stonewort is present. Complete a new form for each pond searched.
- Mark the location: It will be helpful to revisit these areas in future years. To ensure they can be found again by yourself or others please (a) provide an accurate grid reference, or (b) make a sketch of the ponds where you have searched and (c) take lots of photos!

Once completed, enter your results online: <u>www.freshwaterhabitats.org.uk/projects/waternet</u>, or give your recording forms and maps to your regional project officer and we can enter data for you: <u>info@freshwaterhabitats.org.uk</u>.

What it looks like: Bearded Stonewort (a) is a very rare freshwater green algae, known from just a few sites in England. It typically grows in dense clumps with multiple plants (c.30cm in length) growing amongst each other.

It is dark olive to brown in colour and is often encrusted. The most characteristic feature to look out for is the large amount of spine cells on the main stem, giving the plant a bristle-like or 'bearded' appearance (b).

Stoneworts are a difficult group to identify with confidence. To take part in PondNet or to survey Bearded Stonewort at a Flagship Pond

Site https://freshwaterhabitats.org.uk/projects/flagship you will need to attend one of our training sessions.



# Bearded Stonewort (*Chara canescens*) RARE SPECIES RECORDING FORM (PAGE 2 of 4)

| Your name  | Date  |  |
|--|---|--|
| Square: 4 figure grid ref<br>e.g. SP1243 (see your map)  | Pond: 8 figure grid ref<br>e.g. SP 1235 4325 (see your map)   |  |
| Pond name<br>(if known)  |   |  |
| Determiner name ( <u>optional</u> - if<br>someone confirms the identity<br>of the species you've recorded) | Voucher material ( <u>optional</u> -<br>comment if you've taken a photo to<br>confirm identification) |  |

If you find Bearded Stonewort please take a confirmatory photo. You can also take a photo of your pond or your maps (or scan them if you have a scanner) and upload them with the record <u>www.freshwaterhabitats.org.uk/projects/waternet</u>.

## Abundance of Bearded Stonewort in your pond

Record the area of Bearded Stonewort plants from the whole pond, not just the water area, i.e. include areas in the drawdown zone that would be wet in winter, but may be dry in summer. If there are several different patches of Bearded Stonewort in the same pond, use the table below to record the abundance in a small area and add them up - for the analysis **we only need a total**.

| Areas where Bearded Stonewort was found (list): use this table to help with your area calculations, and so you/others can re-find plants on future visits. | Area of Bearded<br>Stonewort (m <sup>2</sup> ) | Area of Bearded<br>Stonewort (%) |
|--|--|----------------------------------|
| 1.   |  |                                  |
| 2.   |  |                                  |
| 3.   |  |                                  |
| 4.   |  |                                  |
| 5.   |  |                                  |

|    | Total area covered by Bearded Stonewort plants (m <sup>2</sup> )<br>Provide a single total for the whole pond based on an actual or estimated area of plants recorded  |
|----|--|
|    | Total area covered by Bearded Stonewort plants (%)<br>Provide a single total for the whole pond based on an actual or estimated area of plants recorded  |
| so | Bearded Stonewort looked for, but not found<br>Note: if you <u>don't</u> find evidence of Bearded Stonewort at the pond, this is an important result so<br>please still enter these findings online (tick box if none found) |

| please still   | enter these findings online (tick box if none found)  |
|--|---|
|  | <u> </u>  |
| Pond sketch map: Make a sketch map of your pond and draw on<br>the location of Bearded Stonewort: use shading if they cover a broad<br>area, or 'x' marks the spot if there are just a few plants. | enter these findings online (tick box if none found)         Location map: Use this box to show the location of the pond and surrounding ponds you searched (or mark the information on the base map included in your site information pack). |
|  | LOTTERY FUNDED  |



| This is a really important part of the survey. Please complete this form whether Bearded Stonewort is present or absent. Each variable provides information known to be linked to site quality and community type, and can be used to investigate reasons for change in Bearded Stonewort occurrence. If you are surveying non-pond habitat – complete all variables that apply.         Go to: www.freshwaterhabitats.org.uk/projects/pondnet/survey-options/habitats       for survey guides and more information (m)         Is the pond new? (less than 10 yrs old) yes, no, unknown       Year of creation? date, decade, unknown       Pond Altitude (m)         Area       Note: This is the surface area of the pond when the water is at its highest level (usually in early spring). It was vegetation like rushes at the pond's outer edge. Measure by pacing (single pace = 0.8-1m) or use online m         Pond dries?       1 = never dries, 2 = Rarely dries: no more than two years in any ten year period, or only drought, 3 = Sometimes dries: dries between three years in ten to most years, 4 = Dries annually. Deduce pond permanence from local knowledge (e.g. landowner) ar personal judgement e.g. water level at the time of the survey. Ponds that dry out annually usually have a hard base.         Overhanging trees & shrubs       This is an estimate of how much of the por directly overhung by trees and shrubs, i.e. would be shaded if the sun was overhead of t   |  |  |  |  |
|---|--|--|--|--|
| Go to: www.freshwaterhabitats.org.uk/projects/pondnet/survey-options/habitats       for survey guides and more informati         Is the pond new? (less than 10 yrs old)       Year of creation?       Pond Altitude (m)         date, decade, unknown       (m)       (m)         Area       Note: This is the surface area of the pond when the water is at its highest level (usually in early spring). It was probably not be the current water level of the pond. The high water level line should be evident from wetlar vegetation like rushes at the pond's outer edge. Measure by pacing (single pace = 0.8-1m) or use online m         Pond dries?       1 = Never dries, 2 = Rarely dries: no more than two years in any ten year period, or only drought, 3 = Sometimes dries: dries between three years in ten to most years, 4 = Dries annually. Deduce pond permanence from local knowledge (e.g. landowner) ar personal judgement e.g. water level at the time of the survey. Ponds that dry out annually usually have a hard base.         Overhanging trees & shrubs       This is an estimate of how much of the por directly overhung by trees and shrubs, i.e. would be shaded if the sun was overhead   |  |  |  |  |
| yes, no, unknown       date, decade, unknown       (m)         Area       Note: This is the surface area of the pond when the water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is at its highest level (usually in early spring). It water is a probably not be the current water level of the pond. The high water level is any teny spring). It water is a spring in the pond of the pond is outer edge. Measure by pacing (single pace = 0.8-1m) or use online material in the pond of the pond of the pond is outer edge. Measure by pacing (single pace = 0.8-1m) or use online material in the pond of the pond is any teny spring. The pond permanence from local knowledge (e.g. landowner) and personal judgement e.g. water level at the time of the survey. Ponds that dry out annually usually have a hard base.         Overhanging trees & shrubs       % of |  |  |  |  |
| Note:       Inis is the surface area of the pond when the water is at its highest level (usually in early spring). It is probably not be the current water level of the pond. The high water level line should be evident from wetlar vegetation like rushes at the pond's outer edge. Measure by pacing (single pace = 0.8-1m) or use online m         Pond dries?       1 = Never dries, 2 = Rarely dries: no more than two years in any ten year period, or only drought, 3 = Sometimes dries: dries between three years in ten to most years, 4 = Dries annually. Deduce pond permanence from local knowledge (e.g. landowner) an personal judgement e.g. water level at the time of the survey. Ponds that dry out annually usually have a hard base.         Overhanging trees & shrubs       This is an estimate of how much of the por directly overhung by trees and shrubs, i.e. would be shaded if the sun was overhead  |  |  |  |  |
| Pond dries?       1 = Never dries, 2 = Rarely dries: no more than two years in any ten year period, or only drought, 3 = Sometimes dries: dries between three years in ten to most years, 4 = Dries annually. Deduce pond permanence from local knowledge (e.g. landowner) an personal judgement e.g. water level at the time of the survey. Ponds that dry out annually usually have a hard base.         Overhanging trees & shrubs       % of pond overhung by trees and shrubs  |  |  |  |  |
| % of pond overhung by trees and shrubs directly overhung by trees and shrubs, i.e. would be shaded if the sun was overhead  |  |  |  |  |
|   |  |  |  |  |
| Waterfowl impact       Major = severe impact of waterfowl e.g. few or no submerged plants, water turbid, pond         1 = major       banks have patches where vegetation removed, feed put down; Minor = waterfowl present         2 = minor       but little impact on pond vegetation, pond still supports submerged plants and banks are no         3 = none       denuded of vegetation; None = no evidence of waterfowl impact (moorhens may be present   |  |  |  |  |
| Fish presence       Major = dense populations of fish known to be present; Minor = small numbers of Crucian         1 = major       Carp, goldfish or stickleback known to be present; Possible = no evidence of fish, but loca         2 = minor       conditions suggest that they may be present; Absent = no records of fish stocking and no revealed during survey.         4 = absent       example   |  |  |  |  |
| Disturbance by dogsMajor = dogs repeatedly use the pond, compacted edges with little vegetation, water very<br>turbid; Minor = dogs use the pond, but little impact on pond vegetation, pond still supports<br>submerged plants and banks are not denuded of vegetation; None = no evidence that dog<br>are using the pond.   |  |  |  |  |
| Aquatic vegetation: includes emergent, floating and submerged plants<br>% of the <u>whole pond</u> (wet and dry) occupied by <u>emergent vegetation</u> – incl.<br>plants like grasses, water mint and rushes, but not floating (e.g. pondweed)<br>%  |  |  |  |  |
| % of pond <u>water surface area</u> covered by all vegetation ( <u>emergent, floating</u><br>( <u>excl. duckweed</u> ) and submerged).  |  |  |  |  |
| Water left in the pond         % of water area in pond relative to maximum water level. This can be 0% if the pond has dried out.         Drawdown. The height drop from the maximum winter water level to current level (see diagram).   |  |  |  |  |
| Grazing         Tick if there is evidence the pond is grazed by livestock. If yes, complete the following boxes:         % of whole pond grazed (note: stock can wade into shallow ponds to graze).         % of pond perimeter grazed (note: stock can wade into shallow ponds to graze otherwise inaccessible edges).         Grazing intensity: rank 1-5 (1=infrequent or low intensity to 5 = margins heavily poached and almost bare).   |  |  |  |  |
| Pond management (tick): use tick boxes to list management within the last 12 months. Use 'other' box for any extra info.         Fully dredged       Partly dredged       >5% vegetation removed       <5% vegetation removed   |  |  |  |  |



# Bearded Stonewort (Chara canescens) **RARE SPECIES RECORDING FORM (PAGE 4 of 4)**

| Water quality:  |  |           |   |  |  |
|---|--|-----------|---|--|--|
| Turbidity / water clarity: Estimate turbidity looking down into c.20cm depth of water in the pond.  |  |           |   |  |  |
| 1 = clear; 2 = moderately clear; 3 = moderately turbid; 4 = turbid  |  |           |   |  |  |
| Inflows and outflows: (tic  | k if infl  | low or ou | tflow present or leave blank)   |  |  |
| Inflow present  |  |           | Dutflow present   |  |  |
| Water chemistry: If suitab  | le kits  | and met   | ers are available (or leave blank)  |  |  |
| рН  |  |           | Conductivity (µS cm-1)  |  |  |
| <b>Nitrate (NO<sup>3</sup>-N ppm):</b> PPW kits provided by FHT <b>Phosphate (PO₄<sup>3-</sup>-P ppm):</b> PPW kits provided by FHT   |  |           |   |  |  |
| (tick one from the following range categories) (tick one from the following range categories)   |  |           |   |  |  |
| <0.2 0.2-0.5 0.5-1  | 1-2  | 2-5       | 5-10 10 + <0.02 0.02-0.05 0.05-0.1 0.1-0.2 0.2-0.5 0.5-1 1 +  |  |  |
|   |  |           |   |  |  |
| Pond base: This refers to the geology (i.e. rock-type) that immediately underlies the pond. You may know, or be able to see the underlying geology in the base or banks of the pond, especially in new ponds. If not, check a geology map or leave this section blank. Choose one of the following to categorise the % composition of each of pond base: 1= 0-32%, 2= 33-66%, 3= 67-100%         Silt/ clay       Sand, gravel, cobbles       Hard rock       Peat       Other (please specify)         Surrounding land use: Estimate the percentage of surrounding land-use in distance zones from the pond perimeter (i.e. the |  |           |   |  |  |
| Habitat   |  | 0-100m    | ond area. In many ponds the 0-5m zone will include surrounding trees/scrub. Examples  |  |  |
| Trees, woodland & scrub   | %  | %         | Deciduous and coniferous woodland, individual trees, scrub and hedgerows.   |  |  |
| Heath & moorland  |  |           | Lowland and upland heathland, moorland and mountain; includes bracken.  |  |  |
| Rank vegetation   |  |           | Unmanaged grass, neglected and abandoned land, set-aside, verges and buffer strips.   |  |  |
| Unimproved grassland  |  |           | Herb-rich, calcareous and acid grassland (good quality plant indicators usually present).<br>Low percentage of agricultural grasses. Not fertilised, little or no drainage. |  |  |
| Semi-improved grassland   |  |           | A transition category. Grasslands modified by fertilisers, drainage, herbicides or intensive grazing, but retaining elements of natural grassland types in the area.        |  |  |
| Improved grassland  |  |           | Fertile agricultural grass, often bright green and lush; including parks and golf greens.   |  |  |
| Arable  |  |           | All crops. Includes flower and fruit crops (e.g. strawberries) and ploughed land.   |  |  |
| Urban buildings & gardens   |  |           | Areas in curtilage (associated with buildings); including glass-houses and farm yards.  |  |  |
| Roads, tracks & paths   |  |           | Including car-parks and footpaths.  |  |  |
| Rock, stone & gravel  |  |           | Cliffs, rock-outcrops, gravel-pits, quarries, areas of sand and gravel or stone.  |  |  |
| Bog, fen, marsh & flush   |  |           | Wetland vegetation and blanket bog.   |  |  |
| Ponds & lakes   |  |           | Permanent and seasonal waterbodies; including trackway pools.   |  |  |
| Streams & ditches   |  |           | Rivers, streams, ditches, springs and canals.   |  |  |
| Other (state)   |  |           | E.g. maritime vegetation, saltmarsh, sand-dune, orchards and railways.  |  |  |
| Is the pond in  | a prot   | ected a   | ea? (e.g. nature reserve, SSSI, etc.) (choose one option - yes, no, unknown)  |  |  |
| New Zealand Pigmyweed   | Cras   | sula hel  | <b>nsii</b> : This non-native weed may have an impact on this species.  |  |  |
| % of drawdown zone occupied by New Zealand Pigmyweed  |  |           |   |  |  |
| Identification of New Zealand Pigmyweed:  |  |           |   |  |  |
| <ul> <li>Can be submerged, err</li> </ul>   | nergent  | and terre | estrial.  |  |  |
| Forms dense mats below and above the water surface.   |  |           |   |  |  |
| <ul> <li>The flowers it has, if an</li> </ul>   | The flowers it has, if any at all, are very small (less than 1cm) whitish- |           |   |  |  |

- Leaves are up to 2cm long in opposite pairs fleshy for emergent plants, but flatter for submerged parts of the plant.
- Similar species (such as the Water-starworts) do not have fleshy leaves. Water-starworts also have • a notch at the leaf tip which is absent in New Zealand Pigmyweed.

#### Other invasive non-native species: (tick all that apply)

Parrot's Feather

Myriophyllum aquaticum

green to slightly pink with 4 petals.

How much of pond perimeter could be surveyed? Note areas of pond not accessible.

Comments box: e.g. new ownership, changes since previous visit, any other information about the pond or survey species.

Floating Pennywort Hydrocotyle ranunculoides

Water Fern Azolla filiculoides Non-native Pondweed, e.g.: Canadian Pondweed Ellodea canadensis, Nuttall's Pondweed Elodea nutallii, Curly Waterweed Lagarosiphon major



